



1. Remove the dressing on postoperative day #1.
2. Apply Band-Aids to small wounds & redress large wound. Please do not use bacitracin or other ointments under the bandage. Use the Cryo/Cuff or ice packs as often as possible for the first 7 days, then as needed for pain relief. An ACE wrap may be used to help control swelling. Do not wrap the ACE too thickly or the cold will not penetrate.
3. You may shower on postoperative days #1-2, however do not wet the wound. Cover the area with tape and plastic wrap or you may also use waterproof Band-Aids. Keep the wound dry at all times, until your sutures are removed by the doctor. Do not soak the knee in water or go swimming until your sutures are removed.
4. Keep your leg elevated with a pillow under your calf, NOT under the knee.
5. Keep the postoperative brace on at all times.
6. Exercises to be performed 2-3 times daily:
  - \* DO NOT bend your knee until directed by Dr. Henshaw.
  - \* Straight leg raises, with foot straight and turned out. Goal is 25 reps 3 times per day.
  - \* Isometric quadriceps contractions.
  - \* Partial weight bearing (up to 50 lbs) while using crutches, with the brace locked in full extension.
7. Please call the office to schedule a follow-up appointment for suture removal 7-10 days after your surgery.
8. If you develop a fever (101.5) or redness or drainage from the surgical incision site, please call our office to arrange for an evaluation.
9. You may experience some low back pain due to muscle spasms from the epidural anesthesia. If so, apply a heating pad to the affected area and take an analgesic if you have not already done so.
10. Enclosed are two prescriptions. Naprosyn, an anti-inflammatory medication, is to be used twice a day **with breakfast and dinner**, for the first 10 days after surgery. Percocet, a narcotic, is to be used on an **as needed** basis for pain in addition to the Naprosyn.

If you have any questions, please feel free to call our office.

If you have any difficulty using anti-inflammatory medications or aspirin, or have a history of ulcer disease, you will only use the Percocet postoperatively.

If you find that the Naprosyn is upsetting your stomach, please call our office to review the medications.